

FIFRA Section 24(c) Special Local Need

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF SOUTH DAKOTA IN THE FOLLOWING GEOGRAPHY: For Control of Weeds in Soybeans in South Dakota West of Highway 281 in the Counties of Aurora, Beadle, Brown, Brule, Buffalo, Campbell, Charles Mix, Douglas, Edmunds, Faulk, Gregory, Hand, Hughes, Hyde, Jerauld, Lyman, McPherson, Potter, Spink, Sully, Tripp, Walworth.

Fomesafen	GROUP	14	HERBICIDE
-----------	-------	-----------	-----------

Sinister®

EPA REG. NO. 5905-593

EPA SLN NO. SD190007

This 24(c) label expires on December 31, 2029

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, and precautions including statements pertaining to the Worker Protection Standards, on the EPA-registered Sinister label.

This label must be in the possession of the user at the time of application.

Failure to follow the directions for use and precautions on this label may result in poor pest control, crop injury, or illegal residues.

Application Methods

Apply SINISTER at 0.49 pt/A as a postemergence broadcast application to soybeans. Apply in a minimum of 15 GPA. Application restricted to ground and aerial applications only.

Spray Additives

Add 1% v/v methylated seed oil (MSO) plus ammonium sulfate (AMS) at a minimum of 8.5 lbs/100 gals of finished spray volume.

Weeds Controlled Weed	Maximum Number of Leaves At Application
Cocklebur, common	2
Mustard, wild	4
Nightshade, black	2
Pigweed, redroot	4
Waterhemp, common	2
Smooth Pigweed	2
Common Ragweed	4, (suppression only)
Palmer Amaranth	2
Common Lambsquarter	2 (suppression only)

Rotational Crop Restrictions Rotational Crops	Minimum Rotational Interval after SINISTER Application (Months)
Bean, Dry Potato Soybean	0 months
Small Grains, such as Wheat, Barley, Rye	9 months
Edible-podded beans and peas not otherwise specified in this label Pea, Dry	12 months
Corn, Field* Corn, Seed*	18 months
All other crops not listed above	Minimum 18 months and a successful field bioassay (refer to Field Bioassay Instructions section) must be performed before planting rotational crop

* For fields with greater than 4% organic matter, a successful field bioassay (refer to **Field Bioassay Instructions** section) must be performed before planting field corn or seed corn.

Field Bioassay Instructions

Plant several strips of the desired crop variety across the field which has been previously treated with SINISTER. Plant the strips perpendicular to the direction SINISTER was applied. The strips should be located so that all the different field conditions are encountered, including differences in soil texture, organic matter, pH, and drainage. If the crop does not show visible symptoms of injury and/or stand reduction, this field can be planted to this crop. If visible injury and/or stand reduction occur, do not plant the field to this crop; plant only a labeled crop.

Use Restrictions

- Do not exceed 0.49 pt/A (0.176 lb ai/A) of SINISTER per acre in any one year. Do not apply to any field more than once every two years.
- Do not make a SINISTER application after June 30.
- Do not harvest soybeans within 45 days of application.
- Avoid overlapping spray swaths, as injury may occur to rotational crops.

SINISTER is a trademark of Helena Holding Company
© Copyright Helena Holding Company, 2025

24(c) registrant:
Helena Agri-Enterprises, LLC
225 Schilling Boulevard, Suite 300
Collierville, TN 38017